

This document is designed to provide information about Laser treatments performed with the **LUTRONIC CLARITY II™** device, their benefits, potential risks, and alternative procedures. It is important that you read this information carefully and have all of your questions answered before signing the consent form.

Most Advanced Laser Platform | All Skin Types | All Year Around | Quick, Comfortable & Effective Treatments | No Downtime

The Technology: FDA approved **LUTRONIC CLARITY II™** is a high-powered dual-wavelength (Nd: YAG 1064nm & Alexandrite 755nm) laser platform offering unmatched capabilities in treating a variety of conditions. Its intelligent features maximize efficacy, comfort, and speed to help achieve desired results. These versatile treatments are safe for all skin types and tones and can be safely used across all areas of the body.

The Procedure: A laser beam produces heat energy to selectively damage independent structures, correcting condition-specific symptoms without harming surrounding tissue. Individual results may vary and are difficult to predict. Noticeable improvements are visible after a single session. Consecutive treatments are highly recommended for optimal results. Future maintenance sessions may be required even if excellent results are achieved. Age, ethnicity, metabolism, medications, supplements (hair, skin, nail, B12, Biotin), hormonal changes, and other factors can affect treatment results. Complete clearance cannot be guaranteed but ideal candidates can achieve significant symptom reduction.

Vascularity & Rosacea: Spider veins under 3mm, broken blood vessels/capillaries, or Telangiectasia Rosacea (blue, purple, or red lines/streaks, webs/branches, or clusters/blotches) Note: veins actively circulate blood and a series of treatments are recommended to achieve a significant long-term reduction in vascularity.

Acne & Cold Sores: Active cold sores (Herpes Simplex Virus - 1 & 2), mild to severe acne breakouts including papules, pustules, nodules, and cystic acne can be effectively treated.

Pigmented Lesions: Primary lesions including sun damage, age spots, freckles, moles, skin tags, macules, papules, patches, and wheals.

Skin Rejuvenation: Improve overall texture, laxity, mild to deep lines and wrinkles, and general skin tightening.

Recommended Course of Treatment: An in-person consultation is recommended to assess suitability for CLARITY II Laser Treatments. Questionable medical conditions and medications will require a clearance from your physician.

Contraindications include, but are not limited to the following:

- Pregnancy or breastfeeding
- Accutane (*isotretinoin*) in the last 6 months
- Retin A topical or other skin-irritating products
- Aesthetic procedures / injections in the last 2 weeks
- Blood-thinning medications within the last week (*Aspirin, iron supplements, herbal supplements such as ginkgo, ginseng, garlic, or St. John's Wort*)
- History of keloid scarring or abnormal wound healing
- Skin cancer or suspicious lesions
- Sunburn (*natural sunlight, tanning*), extreme sensitivity, rosacea
- Poorly controlled medical conditions (*i.e. diabetes, vascular*)
- Impaired immune system and/or use of immunosuppressive medications
- Tattoos and/or permanent makeup
- Implanted medical devices in the treatment area
- Impaired judgment, psychological conditions, and substance or alcohol abuse.
- **Patients with unrealistic expectations**

Risks & Possible Side Effects include, but are not limited to the following:

- Patient failure to follow pre/post care instruction concerning appropriate activity, product use, and sun protection.
- Depending on individual tolerance level, indication, or treatment settings, patients may feel little to slight discomfort during the procedure.
- Common expected skin reactions included erythema (redness), edema (swelling), itching, stinging sensation, and slight irritation or discomfort, which may last 2-4 days. All these reactions may become re-aggravated with heat or sweating.
- Other less common reactions include petechiae (pinpoint scabbing), or purpura (bruising), which can take several days to resolve.
- Activation of individual sensitivities that may include, but are not limited to Herpes Simplex, cold sores, or acne flare-up.
- Any time a skin procedure is performed infection is possible.
- Pigment changes (light or dark spots on the skin) lasting 1-6 months or longer may occur.
- Rarely burns, blisters, or scarring of the treated area may occur.

Although the majority of patients do not experience any side effects and/or complications, you should discuss each of the above with your practitioner to make sure you understand the limitations, risks, potential complications, and consequences of this aesthetic procedure.

Alternative Procedures: CLARITY II Treatments are elective aesthetic procedures that are not medically necessary or required.

Pre / Post Care Instructions: Compliance with the recommended pre / post care instructions as well as recommended follow-up visits by your practitioner are crucial for the success of your treatment and to prevent unnecessary side effects or complications. It is your responsibility to inform your practitioner of any changes in medical history, medications, and most recent sun exposure before each treatment is performed.

Before Your Treatment

- You may not be pregnant or lactating.
- Patients who have used Accutane or similar products within the last 6 months CANNOT be treated.
- Discontinue the use of any topical products that may be drying or irritating (e.g. AHAs, BHAs, retinoids) 3-7 days prior.
- Avoid anti-inflammatory drugs (NSAIDs) for at least 3 days prior. These actions may interfere with the natural inflammatory process critical to proper healing and skin rejuvenation.
- Avoid taking any blood-thinning agents for 3 - 7 days prior unless otherwise prescribed by your physician.
- Avoid unprotected sun exposure, or tanning for at least 2 weeks prior to your procedure. Always use an SPF 30 or higher containing zinc oxide and/or titanium dioxide for proper sun protection.
- Avoid other aesthetic procedures and cosmetic injections in the area(s) to be treated for 2 weeks before and after.
- If you are prone to cold sores, it is recommended to take antiviral medication for 7 days before and after.
- Stay hydrated by drinking 8+ glasses of water daily to improve results & healing time.

Day of Treatment

- Arrive with clean skin free of any lotion, oil, makeup, powder, or sunscreen.
- Notify your practitioner of any changes in your medication, or health status, or personal activities that may be relevant to your treatment.
- Topical numbing using ice packs may be used to maximize your comfort during the treatment.

After Your Treatment

What to Expect:

- Patients will experience mild to moderate erythema (redness), edema (swelling), or a warm stinging / sunburn-like sensation for several days post-treatment, however for more aggressive treatments, these symptoms may last longer than 3 days.
- Other possible skin reactions include irritation, itching, and general sensitivity, which may last for 3 or more days.
- Results are both immediate and progressive. Multiple treatments may be required to achieve individually desired results.
- Commonly patients experience purging of water retention caused by swelling and impurities in the form of water blisters or acne breakouts. This is normal and should resolve within a few days.

48 Hour Care:

- DO NOT wear Makeup for at least 48 hours.
- Avoid all strenuous activity for a minimum of 48 hours as sweating increases infection risk.
- DO NOT USE ICE on the treatment area. DO NOT use any anti-inflammatory drugs (NSAIDs). These actions may interfere with the natural inflammatory process critical to proper healing and skin rejuvenation.
- DO NOT pick, scratch, excessively rub or scrub the treatment area. It is imperative to allow natural healing to minimize post-treatment complications. Treat the skin gently as if you have a sunburn.

General Care:

- Care should be taken to prevent trauma to the treated area especially for the first few days after your procedure.
- You may take Tylenol for post-treatment discomfort or Benadryl / Zyrtec if you experience excessive itching (histamine reaction).
- Avoid All Direct & Indirect Sun Exposure.
- Always use an SPF 30 or higher containing zinc oxide and/or titanium dioxide for proper sun protection. It is safe to begin the use of SPF starting 24-48 hours after your treatment.
- Do not have any other treatments for at least 1 - 2 weeks following treatment or until the skin has completely healed.
- Your practitioner will make individual recommendations on when to resume normal skincare (typically after 2 days).
- Keep the treated area well hydrated (specifically for the first 2 days).

If you experience any side effects, such as prolonged redness or swelling, histamine reaction, blisters, burns, or signs of infection please contact your practitioner immediately for proper assessment and further care instructions.